VISUAL SELECTIVE ATTENTION(U) MEDICAL RESEARCH INST OF SAN FRANCISCO CA K NAKAYAMA 30 MAR 84 AFOSR-TR-84-0774 AFOSR-83-0320

1/1 .

F/G 5/10

AD-A146 220

UNCLASSIFIED



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS -1963 - A



Annual Scientific Report 30 March 1984 AF0SR-83-0320

AD-A146 220

Visual Selective Attention

Medical Research Institute Smith-Kettlewell Institute 2200 Webster Street San Francisco, CA 94115

Dr. Ken Nakayama



FILE COPY

Controlling Office: Air Force Office of Scientific Research/NL Bollling Air Force Base, DC 20332

Approved for public of distribution unlass.

84

٠		REPORT DOCUM	ENTATION PAG	E		
WELASSITIED CATION			1b. RESTRICTIVE MARKINGS			
2L SECURITY CLASSIFICATION AUTHORITY 2L DECLASSIFICATION/DOWNGRADING SCHEDULE 4. PERFORMING ORGANIZATION REPORT NUMBER(S)			Approved for public release; distribution unlimited. E. MONITORING ORGANIZATION REPORT NUMBERIES AFOSR-TR. 34.0771			
se ADDRESS (City, Sinte and ZIP Code) 2200 Webster Street San Francisco, Ca 94115			7b. ADDRESS (City, State and ZIP Code) Bldg # 410 Bolling AFB, DC 20332			
EL NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR -		8b. OFFICE SYMBOL (If applicable) NL	8. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-83-0320			
Bc. ADDRESS (City. State and ZIP Code) Bldg 410 Bolling AFB, DC 20332			10. SOURCE OF FU	NDING NOS.		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NQ.	WORK UNIT
1. TITLE (Include Security Classificati ' VISUAL SELECTIVE AT	-		61102F	2313	A5	1.
12. PERSONAL AUTHOR(S) Nakayama , Ken						
AnnaL	13b. TIME C	OVERED /01/83 to 03/30/8	14. DATE OF REPORT		18. PAGE	COUNT
FIELD GROUP SUB	. GR.	18. SUBJECT TERMS (C	ontinue on reverse if n	ecessary and identi	ify by block numb	er)
This technique has been sinusoidal grating onset recorded on the surface primary visual cortex. to localize the origins to examine the origin of this study will aid it	used to t and of: of the o Because of visual f attent:	differentially fset in differen occipital lobe o current source al evoked potent ton-related pote	localize neurate evidence the riginate in an density analysials, this teentials. It is	at the fiel n area othe sis has suc chnique can s expected	d potential r than the h great abi also be ap that the re	s lity plied sults
20. DISTRIBUTION/AVAILABILITY (UNCLASSIFIED/UNLIMITED 🖾 SA	ME AS RPT.		21. ABSTRACT SEC	SSIFIED	CATION	
Dr. Genevieve M. Haddad			225. TELEPHONE N (Include Area Co 202-767-502	ido) 	22c. OFFICE SY	MBOL
DD FORM 1473, 83 APR		EDITION OF 1 JAN 73	S OSSOLETE.	UNC	ASSIFIED	

84 09 17 015

The Forecast and Progress Report

Selective visual attention has received considerable interest of late. Mounting evidence indicates that the "minds eye" can roam covertly over the visual field at will, lowering thresholds and decreasing reaction time in this region of the visual field (Posner, 1982). Furthermore, it has been hypothesized by Treisman (1980) that the process of selective visual attention may have the important function of synthesizing the disparate features of the visual image which are analyzed in many different visual cortical projection areas (see Allman and Kaas, 1975).

Both single unit electrophysiology in monkeys and massed evoked potential studies in humans show reliable correlates of selective visual attention (Galambos and Hillyard, 1981). The major goal of the present research is to elucidate fundamental mechanisms of visual selective attention by finding an electrophysiological correlate of selective visual attention in monkeys which is comparable to those described in humans. Then we propose to determine the anatomical locus of these changes using the technique of current source density analysis. If successful, such an endeavor will provide an important bridge between human psychophysiological studies of visual attention using evoked potentials and current concepts regarding the anatomical organization of the occipital and parietal cortex in primates.

In order to accomplish this goal, we have purchased a new laboratory computer (PDP 11/23) with funds provided by this research grant. We chose this machine because most of the required applications software for our project has been developed over the past five years at the Max-Planck-Institute in Munich (F.R.G.) for this machine by Mr. Helmut Zucker. We also purchased a Winchester hard disk to have an adequate capacity to store data of many single trials, and the needed interface equipment (clock, ADC, DAC, DIO, etc). Hardware interfaces for the computer have been constructed, tested and debugged. Thanks to a very productive two-week visit by Mr. Zucker in January 1984, these interface functions are now working in conjunction with his software. This includes the smooth working of a graphical output package incorporating a 613 storage oscilloscope coupled to our hardcopy printer.

The main purpose of this software package is to offer to the investigator a specialized programming environment where multichannel analogue data can be sampled, filed efficiently, and easily manipulated by the investigator, without much help from computer specialists. At present we have tested a significant portion of the software package and are now in the process of making the needed modifications so as to make it applicable to our own particular needs, especially in terms of presenting particular blocks of stimulus sequences.

We are also making some technical changes in electrode manufacturing and testing. We have some thin-film miniaturized electrodes manufactured by O. Prohaska in Vienna. These electrodes have been tested in physiological experiments for current source density measurements in the

rabbit visual cortex (Rappelsberger, et al., 1982) and will also be used to examine the origin of visual evoked potentials in monkey cortex (Spekreijse, personal communication). With regards to testing, we are in the process of developing some new in-vivo electrochemical techniques to more conveniently assure the uniformity of electrode characteristics for the duration of chronic implants.

In parallel, we are also conducting experiments with our existing computer and electrode system. Only when the new techniques have been adequately pretested will we replace our present set-up. The proposed experiments for the near-term will concentrate on spatially directed visual attention. This will require that a subject fixate a portion of the visual field, yet forcing a subsidiary awareness at another portion of the visual field by the appropriate behavioral paradigms. These different stimuli are presented in different regions of the visual field and the response to each will be assessed in terms of its proximity to the region of spatially directed attention.

References

Allman, J. and Kaas, J.H. (1975) The dorsalmedial cortical visual area: a third tier area in the occipital lobe of the owl monkey (Aotus frivirgatus). Brain Res. 100, 473-487.

Galambos, R. and Hillyard, S.A. (1981) Electrophysiological approaches to human cognitive processing. Neuroscience Research Program Bulletin 20, 143-265.

Posner, M. (1980) The orienting of attention. Q.J. Exp. Psychol. 32, 3-25.

Rappelsberger, P., Pockberger, H. and Petsche, H. (1982) The contribution of the cortical layers to the generation of the EEG: Field potential and current source density analysis in the rabbits visual cortex. EEG and Clin. Neurophysiol. 53, 254-269.

Treisman, A. (1982) Perceptual grouping and attention in visual search for features and for objects. J. Exp. Psychol: Human Perception and Performance 8, 194-214.

